
5-1-2003

The Mayfly Newsletter

Peter M. Grant

Southwestern Oklahoma State University, peter7grant@gmail.com

Follow this and additional works at: <https://dc.swosu.edu/mayfly>




Part of the [Biology Commons](#), [Entomology Commons](#), [Systems Biology Commons](#), and the [Zoology Commons](#)

Recommended Citation

Grant, Peter M. (2003) "The Mayfly Newsletter," *The Mayfly Newsletter*. Vol. 12: Iss. 2, Article 1.
Available at: <https://dc.swosu.edu/mayfly/vol12/iss2/1>

This Article is brought to you for free and open access by the Current Newsletters at SWOSU Digital Commons. It has been accepted for inclusion in The Mayfly Newsletter by an authorized editor of SWOSU Digital Commons. An ADA compliant document is available upon request. For more information, please contact phillip.fitzsimmons@swosu.edu.



THE MAYFLY NEWSLETTER

Vol. 12 No. 2

Southwestern Oklahoma State University, Weatherford, Oklahoma 73096-3098 USA

May 2003

Mayfly Diversity in the Headwaters of Brazilian Altitudinal Streams (Serra do Cipó National Park)

Michael D. C. Goulart, supervised by Dr. Marcos Callisto,
Lab. of Benthic Ecology, Department of General Biology, Institute of Biological Sciences, Federal University of Minas Gerais, Belo Horizonte, MG, Brazil
michaeld_03@hotmail.com

Introduction

Serra do Cipó is located in the central part of the Minas Gerais State (19°-20° S; 43°-44° W), the southeastern region of Espinhaço Mountain Chain. The vegetation is composed of savanna (locally called "cerrado") in the lower altitudes (up to 1000 m), rupestrian fields in the highest portions (above 1000 m) and riparian forest in the humid valleys along the rivers (Giulietti, 2002). The climate is classified as Cwb (Köppen), with rainy summers and dry winters and an annual median of pluviosity about 1500 mm (Galvão & Nimer, 1965).

The Ephemeroptera are a conspicuous and diverse group in the sediment and on aquatic macrophytes in lotic ecosystems of Serra do Cipó, and often constitutes the dominant group in the macroinvertebrate communities (Galdean *et al.*, 1999, 2000, 2001). Within these ecosystems, some are located in nearly "pristine" areas (e.g., Indaiá and Capão da Mata streams, located within the Serra do Cipó National Park), while others are located in regions that have been affected by human activities to varying degrees (Cipó, Peixe and Preto do Itambé rivers) (Galdean *et al.*, 2000).

The aims of my master dissertation were first to evaluate the distribution, diversity, structure and substrate associations of mayfly assemblages in these five representative lotic ecosystems of Serra do Cipó (Brazil), determining the major factors influencing the mayfly assemblages diversity and structure. Second, I studied some of the premises of the river continuum concept trying to evaluate aspects of the longitudinal distribution, diversity patterns and substrate association of mayfly nymphs in the longitudinal gradient of Indaiá stream, located inside the Serra do Cipó National Park.

Methods

The samples were collected during the rainy and dry periods of 1998, 2000 and 2001, using kick nets (0.250 mm mesh, 1m² of sampled area), and Surber sampler (0.250 mm mesh, 0.01 m² and 0.0625 m²) and preserved with 10% formalin. In the laboratory, the samples were washed through a 0.250 mm sieve, sorted and identified with a

stereomicroscope, and the identified organisms were preserved in 70% alcohol and deposited in the Reference Collection of Benthic Macroinvertebrates, Institute of Biological Sciences, Federal University of Minas Gerais (Callisto *et al.*, 1998). The physical and chemical parameters of the water column were measured "in situ" using a Horiba multiprobe (temperature, pH, electrical conductivity and dissolved oxygen). The analysis of the total alkalinity and the concentrations of total and dissolved nutrients were performed in the laboratory (Golterman *et al.*, 1978; Mackereth *et al.*, 1978; Carmouze, 1994).

Results Summary and Discussion

Since the preliminary studies in Serra do Cipó, accomplished by Galdean *et al.* (1999, 2000, 2001), taxonomic identification of several mayfly genera were added or corrected by using updated international literature (e.g., Salles, 2002; Dominguez *et al.*, 2001). At that time, 19 genera, within 5 families, were identified by Prof. Galdean. The dominant genera found by these authors were *Baetis* (Baetidae) and *Farrodes* (Leptophlebiidae). Besides their efficiency to explore the main available trophic resources of the investigated ecosystems, they also showed a larger amplitude to adverse abiotic conditions, such as higher nutrient concentrations and fine sediment granulometry. With the increasing taxonomic revisions and further collections in South America, developed by researchers such as Drs. C. Lugo-Ortiz, W. P. McCafferty, E. Dominguez,

Continued on p. 2

Inside This Issue

History of Insects, IJM 2004, WWW	3
Books About Mayflies	4
Baetidae of Madagascar	5
Conference Guidelines, Postdoc Request, Publication Info, National Geographic Article	6
2002 Mayfly Bibliography	7
Cartoon	15
Address Update	16



Continued from p. 1

MSc. C. Molineri, and MSc. F. Salles, various new taxa were described and/or classified under different nomination. For instance, the genus *Baetis* is now recognized only in a North and Central American distribution. In this way, the former specimens identified under the old classification (*Baetis*) are now being classified as *Americabaetis* and *Cryptonympha*.

In total, 30 genera, distributed within 7 families, were identified, with a high dominance of Baetidae (mainly *Paracloeodes* spp. and *Americabaetis* spp.), Leptohiphidae (mainly *Leptohiphes* spp., although, some revision on the collected species are needed) and Leptophlebiidae (mainly *Askola* cf. *froehlichii* Peters, 1969 and *Miroculis* sp.). My study shows that most of the identified taxa present specific substrate associations, being found in 2-3 substrates. Some genera, besides being restricted found in association with few substrates, also show a clinal pattern in their longitudinal distribution, such as *Campylocia* cf. *bocainenses* (Euthyplociidae), found only in gravel/sand deposits in shallow and low flow areas, from 2nd to 4th order. It could also be observed the probable influence of temperature in the distribution of Ephemeroptera. Galdean *et al.* (1999) points out the possible competition between the Baetidae and Leptophlebiidae family in Serra do Cipó lotic ecosystems. I found that in the headwaters, the mayfly assemblages were composed mostly of leptophlebiid genera, with high diversity and abundance. Otherwise, in the lower stretches of the longitudinal gradient (e.g., 4th order and further), where temperatures reached higher values, we observed a modification in the assemblages structure, with the dominance of the baetid family presenting high numbers and higher diversity.

The main results that I obtained, besides the increased taxonomic survey of Serra do Cipó lotic ecosystems, include new distributional records of recently described mayfly genera and species and ecological data about little known Ephemeroptera taxa (*Paracloeodes*; *Americabaetis*; *Aturbina georgei* Lugo-Ortiz & McCafferty, 1996; *Apobaetis fiuzai* Salles & Lugo-Ortiz, 2002 - Baetidae; *Tricorythopsis* - Leptohiphidae; *Campylocia* cf. *bocainenses* Pereira & da Silva, 1990). This information, together with the assessment of ecological characteristics and preservation status of river stretches, contributes to the knowledge about the main factors governing the distribution and diversity of mayfly assemblages. These patterns are being used as tools for conservation strategies of altitudinal headwaters streams in Serra do Cipó (MG - Brazil).

Final Considerations

Some researchers (e.g., E. Dominguez, C., Molineri, F. Salles, P. Grant, L. Corkum, D. Hilsenhoff, D. Dudgeon, M. Sartori, J. Alba-Tecedor, C. Hawkins, N. Kluge, I. Campbell, J. Brittain and several others) helped a lot sending me reprints of papers about mayflies and/or benthic macroinvertebrate ecology or checking some material. However, there are still great difficulties in finding appropriate and recent literature about taxonomy and ecology of mayflies. I hope that our studies in the headwaters of Serra

do Cipó can help the scientific progress of mayfly ecology to elucidate some of the remaining questions in this aquatic insect order research in the Neotropics. In the future (probably during my Ph.D. thesis), it will be necessary to expend greater efforts on species identification. Future collaborations will be welcome.

References Cited

- CALLISTO, M.; BARBOSA, F. A. R. & VIANNA, J. A. 1998. Qual a importância de uma coleção científica de organismos aquáticos em um projeto de biodiversidade? *Anais do IV Simpósio de Ecossistemas Brasileiros*. 2(1): 432-439.
- CARMOUZE, J. P. 1994. O metabolismo dos ecossistemas aquáticos: fundamentos teóricos, métodos de estudo e análises químicas. Ed. Edgard Blücher: FAPESP. 253p.
- DOMÍNGUEZ, E.; HUBBARD, M. D.; PESCADOR, M. L. & MOLINERI, C. 2001. Ephemeroptera. In: H. R. Fernandez & E. Dominguez (eds.) *Guía para la determinación de los artrópodos bentónicos sudamericanos*. pp. 17-54. Ed. Universidad Nacional de Tucumán, Argentina.
- GALDEAN, N., BARBOSA, F., CALLISTO, M., ROCHA, L., MARQUES, M. 1999. A proposed typology for the rivers of Serra do Cipó (Minas Gerais, Brazil) based on the diversity of benthic macroinvertebrates and the existing habitats. *Travaux Museum History Natural Grigore Antipa*. 41 (1): 445-453.
- GALDEAN, N., CALLISTO, M., BARBOSA, F. 2000. Lotic ecosystems of Serra do Cipó, southeast Brazil: water quality and a tentative classification based on the benthic macroinvertebrate community. *Aquatic Ecosystem Health and Management*, in press.
- GALDEAN, N., CALLISTO, M., BARBOSA, F. 2001. Biodiversity assessment of benthic macroinvertebrates in altitudinal lotic ecosystems of Serra do Cipó (MG-Brazil). *Brazilian Journal of Biology*, 61 (2): 239-248.
- GALVÃO, M. V. & NIMER, E. 1965. Clima. *Geografia do Brasil-Grande Região Leste* 5: 91-139, IBGE, Rio de Janeiro.
- GIULIETTI, A. M. 2002. Flora: diversidade, distribuição geográfica e endemismos. *Serra do Cipó: ecologia e evolução* (ed. G. W. Fernandes), in press.
- GOLTERMAN, H. L.; CLYMO, R. S. & OHNSTAD, M. A. M. 1978. *Methods for chemical analysis of freshwaters*. Blackwell Scientific Publications, 210 p.
- LUGO-ORTIZ, C. R. & MCCAFFERTY, W. P. 1996. *Aturbina georgei* gen. et sp. n.: a small minnow mayfly (Ephemeroptera: Baetidae) without turbinate eyes. *Aquatic Insects*. 18: 175-183.
- MACKERETH, F. J. H.; HERON, J. & TALLING, J.F. 1978. *Water Analysis: some revised methods for limnologists*. Freshwater Biological Association. Windermere 120 p.
- PEREIRA, S. M. & DA SILVA, E. R. 1990. Nova espécie de *Campylocia* Needham & Murphy, 1924 com notas biológicas (Ephemeroptera, Euthyplociidae). *Buletim*

Continued on p. 3

History of Insects

Dr. Nikita Julievich Kluge, Department of Entomology, Biological Faculty, S.-Petersburg State University, 199034 Russia, <http://www.bio.pu.ru/win/entomol/kluge-en.htm>

A new book, *History of Insects*, edited by A.P. Rasnitsyn and D.L.J. Quicke, has been published by Kluwer Academic Publishers (P.O. Box 322, 3300 AH Dordrecht, The Netherlands, or 101 Philip Drive, Norwell, MA 02061, U.S.A.). It was prepared as the second edition of the book, *Historical Development of Class Insects*, published in Russian in 1980 in the series "Trudy Paleontol. Inst. Akad. Nauk SSSR," vol. 175, edited by B.B. Rohdendorf and A.P. Rasnitsyn. In fact, the new version is strongly changed and is several times thicker (517 large pages instead of 268 smaller ones).

"This is the first time that a single book has attempted to cover the whole fossil history of insects so comprehensively. The volume embraces the history of insect paleontology, the methods of studying fossils, the taphonomic processes leading to their formation, the diagnostic features of all insect orders, both extant and extinct, the major fossils of each order, and the implications that can be drawn from the paleontological record about past ecology and climates.

"Many new insights are presented. It is the product principally of the largest paleontological group in the world, in Moscow, and makes full use of the remarkable collection that these workers have developed. It includes a very large number of illustrations showing both real fossils and reconstructions of extinct taxa.

"The systematic part is treated in a phylogenetic framework with information on fossil groups being used to help interpret relationships. An appendix provides information on virtually all sites where fossil insects have been found.

"This book is essential to all students of paleontology and contains a wealth of information that will be of interest to students of evolutionary relationships and of paleontology in general."

The book is built like the Tower of Babel: there are 22 co-authors, each with their own opinion about insect phylogeny, thus the text is full of contradicting statements. Particularly, the characteristics of mayflies in the narrower sense ["2.2.1.1.1.3. ORDER EPHEMERIDA Latreille, 1810. THE TRUE MAYFLIES (=Ephemeroptera Hyatt et Arms, 1891 (s.l.); =Euephemeroptera Kluge, 2000)"] is written by N.Ju. Kluge and N.D. Sinitshenkova, who could not agree with the characteristics of mayflies in a wider sense written by A.P. Rasnitsyn ["2.2.1.1.1. SUPERORDER EPHEMERIDEA

Latreille, 1810. THE MAYFLIES (=Panephemeroptera Crampton, 1928)"].

An unfortunate misprint is Fig. 84 which presents an old Demoulin's (1956) reconstruction of *Triplosoba pulchella* and the legend stating that this is "restored by N.Yu.Kluge (orig., based on Brongniart 1893 and Carpenter 1963a)."

Nevertheless, the book really contains a lot of important information on the fossil history of insects, being the most comprehensive source on this subject. Analysis of phylogeny of neopteran groups given in this book is most comprehensive this time, until the continuation of the book, *Modern Systematics of Insects* (Kluge, 2000) which is not finished.

2004 International Joint Meeting

XI International Conference on Ephemeroptera
XV International Symposium on Plecoptera

Flathead Lake Biological Station and
Division of Biological Sciences
The University of Montana
Polson, Montana (USA)
August 22 - 28, 2004

For further information or to be added to the mailing list, send an email to Ms. Sue Gillespie, sgill@selway.umt.edu or go to the web page at <http://www.umt.edu/biology/flbs/Events/StoneMay2004.htm>.

Continued from p. 2

do Museu Nacional, N.S., Rio de Janeiro, Zoologia, No. 336:1-12.

PETERS, W. L. 1969. *Askola froehlichii*, a new genus and species from southern Brazil (Leptophlebiidae: Ephemeroptera). Florida Entomologist 52:253-258.

SALLES, F. F. 2002. *Contribuição ao estudo da taxonomia da família Baetidae (Insecta: Ephemeroptera) na região sudeste do Brasil*. Master dissertation. Universidade Federal de Viçosa, Viçosa, Minas Gerais. 129 p.

SALLES F. F. & LUGO-ORTIZ C. R. 2002. A distinctive new species of *Apobaetis* (Ephemeroptera: Baetidae) from Mato Grosso and Minas Gerais, Brazil. *Zootaxa*. 35, 1-6.

Mayfly WWW Sites

Mayfly Central

<http://www.entm.purdue.edu/entomology/research/mayfly/mayfly.html>

Ephemeroptera Galactica

<http://www.famu.org/mayfly/>

Entomologie für Fliegenfischer

Walter Reisinger, Ernst Bauernfeind, Erhard Loidl. 2002. Vom Vorbild zur Nachahmung. Ca. 288 Seiten, ca. 500 Fotos. Gebunden. Format 19 cm x 24,5 cm. ISBN 3-8001-3594-9.

In the last issue of this newsletter, I announced the availability of a new book, *Entomologie für Fliegenfischer*, by Reisinger et al. I was fortunate to receive a copy of this book and must comment on it further.

This book is divided into four sections. Each of the first three sections focuses on mayflies, stoneflies, or caddisflies. The sections begin with general information about the biology of the order. Following that is information about species of interest to fly fishing, organized by family. Each section ends with a table with flight period information for the species discussed.

The last section contains information about fly tying

What really caught my eye were the color photographs - over 500. There are photographs of habitat, the adult stage of the species discussed, and tied flies. These photographs are exquisite. They are very clear and show much detail.

For further information, contact the publisher at Verlag Eugen Ulmer GmbH & Co., Postfach 70 05 61, D-70574 Stuttgart, phone (0711) 4507-0, fax (0711) 4507-120, email info@ulmer.de.

The web site describing this book in more detail is located at <http://www.ulmer.de/Vorlagen/Webapp/ULMERSHOP/Load/AllgemeinTexte/veu.htm>.

Proceedings of the Fourth International Conference on Ephemeroptera

V. Landa, T. Soldan and M. Tonner, eds. 1984.

Three copies of these proceedings remain, which we believe to be the last copies available. To purchase a copy, send US\$50.00 to Peter Grant, Department of Biological Sciences, Southwestern Oklahoma State University, 100 Campus Drive, Weatherford, Oklahoma 73096-3098, USA. To reserve a copy, you might want to email me first at grantp@swosu.edu.

Money earned from the sale of these proceedings will be deposited in the William L. Peters Scholarship Fund.

Trends in Research in Ephemeroptera and Plecoptera

Eduardo Dominguez, ed. 2001. Kluwer Academic/Plenum Publishers. ISBN 0-306-46544-2. 490 pp. EUR 156.00, USD 135.00, GBP 95.00.

These are the proceedings of the IXth International Conference on Ephemeroptera and the XIIIth International Symposium on Plecoptera, held in Tucuman, Argentina, in 1998.

After much anticipation, this book has been published and is now available for purchase. Conference participants should have received their personal copies in the mail.

See the publisher's website (<http://www.wkap.nl/prod/b/0-306-46544-2?a=2>) for additional information or to order a copy.

Books About Mayflies

Phylogenetic System of Ephemeroptera. Vol. 1

Dr. Nikita Julievich Kluge, Department of Entomology, Biological Faculty, S.-Petersburg State University, 199034 Russia, <http://www.bio.pu.ru/win/entomol/kluge-en.htm>

This book is being prepared for publication on paper. Its first version, *Draft Revision of Supraspecies Taxa of Ephemeroptera (Without Atalophlebiinae)*, has been available in English text since 1988. The second version, *Revision of Supraspecies Taxa of Ephemeroptera (Except for Atalophlebia/fg1)*, has been available in Russian text since 2000.


The new text markedly differs from the previous ones. Here are presented only the contents, classificational scheme, foreword and indexes of taxa names.

Look please to the index of species names and the index of supraspecies taxa names to search for mayfly species or genera which you or your colleagues introduced or changed in status (apart from those belonging to Baetidae and Leptophlebiidae). Some information about it can be found on the web at <http://www.bio.pu.ru/win/entomol/KLUGE/EPH/Contents.htm>.

It is not excluded that I missed something important in recent publications, so I would be thankful to everybody to inform me about this by email (kluge@ent.bio.pu.ru).

Etude Systématique, Cladistique et Biogéographique des Baetidae (Ephemeroptera) de Madagascar

Systematic, Cladistic and Biogeographical Study of the Malagasy Baetidae (Ephemeroptera)



Jean-Luc Gattolliat

Musée Cantonal de Zoologie, Lausanne & Université de Lausanne,
Faculté des Sciences,
Institut d'Ecologie, SWITZERLAND
jean-luc.gattolliat@serac.vd.ch

Mayflies (Ephemeroptera) are among the oldest known flying insects and encompass a very small number of species (ca. 2500 species). Larvae are strictly freshwater inhabitants; this stage lasts generally one year. The imaginal stage is extremely short, from a few hours to a few days, and is devoted almost entirely to reproduction.

Madagascar is the fourth largest island in the world by area. It is situated in the western part of the Indian Ocean, at a distance of more than 300 km from the African coast. Madagascar belonged to Gondwana. It was first separated from the African plate (-165 M.y.), then moved to the South (-65 M.y.), before the break-off with the Indian plate (-65 M.y.).

Knowledge of the Malagasy mayflies was until recently extremely poor. The program *Biodiversity and Biotypology of Malagasy Freshwaters*, jointly run by the French ORSTOM and the Malagasy CNRE, began a global survey of the freshwater macroinvertebrates. The systematics of several mayfly families (Tricorythidae, Polymitarcyidae, Palingeniidae,...), and other invertebrate groups (caddisflies, blackflies,...) was the subject of ground studies. Our present study is one part of this global program.

Until the middle of the nineties, only four baetid species belonging to three different genera had been described from Madagascar. During the last six years, 25 papers were dedicated to the systematics of the Baetidae, allowing the description of 50 new species and 8 new genera. The Malagasy fauna now encompasses 22 genera and 54 species. Despite its size, Madagascar has the same diversity, at specific and generic levels, as a continent.

Our knowledge of the Baetidae is sufficient to perform a cladistic and biogeographical study. Our phylogenetic reconstruction allows us to propose five main lineages and to indicate, for each of them, the genera included and their features.

The Malagasy Baetidae possess a high level of endemism: 53 of the 54 species and one third of the genera are endemic. It shows extremely strong affinities with the African fauna, as more than 90% of the genera present in Madagascar or in Africa have a distribution restricted to this area. Other components, especially Oriental and Oceanian, are negligible. These areas share with Madagascar only a few widespread genera. These African affinities are in contradiction with the geological events, especially the break-off history of Gondwana.

Some explanations can be given to solve this contradiction. The most likely is that the dispersal power of the mayflies, especially of the Baetidae, is greatly underestimated. The study of recent volcanic islands, particularly of the Comoros, clearly demonstrates that the Baetidae are able to disperse over more than 300 km. Consequently, a colonisation by the Baetidae of Madagascar from continental Africa, after the break-off, must be considered as possible.

We have established scenarios explaining the biogeographical history of each of the five lineages. For four of them, Africa has to be regarded as the centre of origin. The fifth lineage probably has a Palearctic origin; Africa should be considered as a secondary centre of speciation. These lineages should have secondarily colonised Madagascar from continental Africa.

This work opens up new perspectives. It allows the use of the Baetidae for faunistic and ecological studies, especially for problems related to water quality. It must be also considered as a first step for understanding the dispersion and colonisation of the islands of the western part of the Indian Ocean.

[Editor's note: Jean-Luc completed his Ph.D. thesis last spring on the above research. A PDF version of his thesis is available on CD. Contact him at the above address for a copy.]

Request for Postdoctoral Position

My doctoral dissertation, which is now finished, had three components: (1) reconstruction of the phylogeny of Ephemeroptera; (2) classification of the Chinese Ephemeroptera, and (3) the relationship between mayflies and water quality.

I am very interested in mayflies and in using them to monitor water quality, especially of rivers. So, I hope to continue my studies in the USA someday. With the developing of society, the pollution of water is serious, and there are few researchers in this field in China.

So, I would like to locate a postdoctoral position on mayflies or aquatic biology. Please contact me if you might have some information regarding my request: ZHOU Changfa, Department of Biology, Nankai University, Tianjin, 300071, Peoples Republic of China, email cfzhou@eyou.com.

Mayfly Newsletter Update

- Due to an error on some mailing labels, a number of February 2002 newsletters were returned, particularly those sent overseas. If you did not receive a copy of the February 2002 newsletter, please let me know and I will mail one to you.
- The last issue of *The Mayfly Newsletter* was February 2002, 12(1).
- A new mailing label format is now in use. Please check your mailing label for accuracy. Let me know if changes need to be made.



National Geographic

Check out the article on *Palingenia longicauda* in the May 2003 issue of *National Geographic*. The photographs are gorgeous!

Guidelines for Submitting Proposals to Host an International Conference

As reported in the February 2002 newsletter, the representatives from the International Conferences on Ephemeroptera and the International Society of Plecopterologists agreed that there should be a set of guidelines for submitting proposals to host conferences. This was decided during the joint business meeting on 9 August 2001 in Perugia, Italy. Peter Zwick, Michel Sartori, and Peter Grant were appointed to prepare the following guidelines.

Preliminary Proposals

Preliminary proposals to host a conference may be submitted six years (i.e., two conferences) prior to the year of the proposed conference, but a final vote on the conference site will not be made until three years (i.e., the

previous conference) prior to the actual conference date.

Final Proposals

1. Proposals should be submitted at least one month prior to the conference during which the proposal will be officially presented.
2. A copy of this proposal should be sent to the chair of each committee - International Conferences on Ephemeroptera and the International Society of Plecopterologists.
3. Proposals should be submitted by email. This facilitates distribution of the proposal to the members of the two committees.
4. Proposals should contain detailed information regarding plans to host the conference.

Eaton's Revisional Monograph for Sale

Bob Boyle reports that he has parts 2 and 3 of Eaton's "A Revisional Monograph of Recent Ephemeridae or Mayflies" (*The Transactions of the Linnean Society of London*): Part 2, July 1884, pages 77-152 of text, 20 pages of plates (Plates 25-45); Part 3, April 1885, pages 153-230 of text, Plates 46-63. Description: small folio, original brown paper covers, covers slightly age worn, some text pages unopened.

He is asking for US\$200, which is what he paid a British dealer some time ago.

Contact Bob at Shad Roe, Lane Gate Road, Cold Spring, New York 10516 USA, phone (845) 265-2944, fax (845) 265-7838.

Proceedings of the First Symposium of Aquatic Entomologists

The proceedings of the first symposium of aquatic entomologists in East Asia has recently been published:

Bae, Y. J., ed. 2001. *The 21st Century and Aquatic Entomology in East Asia. Proceedings of the 1st Symposium of Aquatic Entomologists in East Asia.* The Korean Society of Aquatic Entomology, Korea.

These proceedings contain 10 papers (3 on mayflies) and one invited lecture.

Contact Dr. Bae for more information: Y. J. Bae, Biology, Seoul Womans University, 126 Kongnung 2-Dong, Nowon-Gu Seoul 139-774, KOREA.

2001 Mayfly Bibliography

[Editor's note: This bibliography was published as the Ephemeroptera portion of the 2001 (2002) North American Benthological Society's (NABS) *Current and Selected Bibliography on Benthic Biology*.]

The following is a list of current publications on Ephemeroptera that have been published up to and during 2001 and have not appeared in previous NABS Bibliographies.

I would appreciate receiving a reprint or complete bibliographic reference of any article about mayflies, especially if it contains scientific names, so that it may be included in next year's bibliography. Also, I would like to be informed of any corrections or omissions in this or past bibliographies. Suggestions are always welcome.

Please send all correspondence to Peter M. Grant, Department of Biological Sciences, Southwestern Oklahoma State University, 100 Campus Drive, Weatherford, Oklahoma 73096-3098 USA, phone (580) 774-3294, fax (580) 774-7140, email grantp@swosu.edu.

If you would like an electronic copy of this year's mayfly bibliography, simply send me a request via email. I will send this file to you as an attachment. This bibliography is also available on my website: <http://www.swosu.edu/~grantp/research.htm>.

- Adamek, Z.; Sukop, I. 2001. The role of supplementary feeding in food competition between common carp (*Cyprinus carpio*) and perch (*Perca fluviatilis*) in a pond culture. *Krmiva* 43: 175-184.
- Alba-Tercedor, J.; Picazo-Munoz, J.; Jaimez-Cuellar, P. 2000. Presence of *Labiobaetis neglectus* (Navas, 1913) (Ephemeroptera, Baetidae) in the south of the Iberian Peninsula. *Bol. Asoc. Espanola Entomol.* 24(3-4): 221-222. (In Spanish)
- Alba-Tercedor, J.; Picazo-Munoz, J.; Jaimez-Cuellar, P. 2000. First record of *Pseudocentropilum nanum* (Bogoescu, 1951) (Ephemeroptera, Baetidae) in the Iberian Peninsula. *Bol. Asoc. Espanola Entomol.* 24(3-4): 224. (In Spanish)
- Alba-Tercedor, J.; Puig-Garcia, M. A. 2000. *Labiobaetis tricolor* (Tshernova, 1928) (Ephemeroptera: Baetidae) misrecorded in the Iberian Peninsula. *Bol. Asoc. Espanola Entomol.* 24(3-4): 223. (In Spanish)
- Amundsen, P. A.; Gabler, H. M.; Ruse, L. S. 2001. Intraspecific food resource partitioning in Atlantic salmon (*Salmo salar*) parr in a subarctic river. *Aquat. Living Resour.* 14(4): 257-265.
- Aranha, J. M. R.; Gomes, J. H. C.; Fogaca, F. N. O. 2000. Feeding of two sympatric species of *Characidium*, *C. lanei* and *C. pterostictum* (Characidiinae) in a coastal stream of Atlantic Forest (southern Brazil). *Brazilian Arch. Biol. Tech.* 43(5): 527-531.
- Arnold, G.; Smith, C.; Cobb, D. G.; Kidd, K. A.; Hesslein, R. H.; Lockhart, W. L. 1997. Mayflies as biomonitors of mercury and cadmium in the Lake Winnipeg basin. *Canadian Tech. Rep. Fish. Aquat. Sci.* 2163: 51-52. [Abstract]
- Bacescu, M. 1997. Homage to Professor Constatin Motas: First contacts and collaborations with Professor Constatin Motas. *Trav. Mus. Nat. Hist. Natur. "Grigore Antipa"* 37: 330-332. (In French)
- Bae, Y. J.; Lee, B. H. 2001. Human impacts on stream ecosystems and freshwater arthropods in Korea. *Korean J. Entomol.* 31(2): 63-76. (In Korean; English summary)
- Bae, Y. J.; Liu, G. C. 1999. Mayflies (Ephemeroptera) from Changbaishan area in Northeast Asia. *Entomol. Res. Bull.* 25: 1-6.
- Ball, S. L. 2000. Evolutionary ecology and population genetics of tytoparthenogenesis in the mayfly, *Stenonema femoratum* (Ephemeroptera: Heptageniidae). Dissertation, University of Missouri – Columbia.
- Baptista, D. F.; Buss, D. F.; Dorville, L. F. M.; Nessimian, J. L. 2001. Diversity and habitat preference of aquatic insects along the longitudinal gradient of the Macae River basin, Rio de Janeiro, Brazil. *Brazilian J. Biol.* 61(2): 249-258.
- Baptista, D. F.; Dorville, L. F. M.; Buss, D. F.; Nessimian, J. L. 2001. Spatial and temporal organization of aquatic insect assemblages in the longitudinal gradient of a tropical river. *Brazilian J. Biol.* 61(2): 295-304.
- Barmuta, L. A.; McKenny, C. E. A.; Swain, R. 2001. The responses of a lotic mayfly *Nousia* sp. (Ephemeroptera: Leptophlebiidae) to moving water and light of different wavelengths. *Freshwater Biol.* 46(5): 567-573.
- Bass, D. 2000. A preliminary study of aquatic macroinvertebrates from two springs in the Pontotoc Ridge Nature Preserve, Oklahoma. *Proc. Oklahoma Acad. Sci.* 80: 105-109.
- Bauernfeind, E.; Humpesch, U. H. 2001. Mayflies of Central Europe – identification and ecology (Insecta: Ephemeroptera). *Naturhistorischen Mus. Wien.* (In Austrian; English summary)
- Bauernfeind, E.; Moog, O. 2001. Comments on the systematics and nomenclature of selected Austrian mayflies (Insecta: Ephemeroptera). *Lauterbornia* 40: 79-91.
- Baumann, R. W.; Kondratieff, B. C. 2000. A confirmed record of the Ephemeroptera genus *Baetisca* from west of the continental divide and an annotated list of the mayflies of the Humboldt River, Nevada. *West. North Amer. Natur.* 60(4): 459-461.
- Bello, C.; Carlos, L. 2000. Taxonomic descriptions of the nymphs of six genera of the family Leptophlebiidae (Insecta: Ephemeroptera) from Paso del Diablo Creek, Guasare Coal Mine Region (Zulia State). *Ciencia* 8(2):

- 127-136. (In Spanish, English abstract)
- Benke, A. C.; Wallace, J. B.; Harrison, J. W.; Koebel, J. W. 2001. Food web quantification using secondary production analysis: Predaceous invertebrates of the snag habitat in a subtropical river. *Freshwater Biol.* 46(3): 329-346.
- Benstead, J. P.; Barnes, K. H.; Pringle, C. M. 2001. Diet, activity patterns, foraging movement and responses to deforestation of the aquatic tenrec *Limnogale mergulus* (Lipotyphla: Tenrecidae) in eastern Madagascar. *J. Zool.* 254(1): 119-129.
- Berrady, I.; Essafi, K.; Mathieu, J. 2000. Comparative physico-chemical and faunal studies of two thermal springbrooks near Sidi Harazem (Morocco). *Ann. Limnol.* 36(4): 261-274.
- Berthold, E.; Egge, C.; Schuller, I. 2001. First find of *Metreletus balcanicus* (Insecta, Ephemeroptera, Ameletidae) in Bavaria. *Lauterbornia* 40: 93-97. (In German; English abstract)
- Besser, J. M.; Brumbaugh, W. G.; May, T. W.; Church, S. E.; Kimball, B. A. 2001. Bioavailability of metals in stream food webs and hazards to brook trout (*Salvelinus fontinalis*) in the upper Animas River watershed, Colorado. *Arch. Environ. Contam. Toxicol.* 40(1): 48-59.
- Boiteau, G.; Osborn, W. P. L.; Xiong, X.; Bousquet, Y. 2000. The stability of vertical distribution profiles of insects in air layers near the ground. *Can. J. Zool.* 78(12): 2167-2173.
- Bonali, F.; Impieri, A. 2000. Signaling of *Potamanthus luteus* L. (Insecta-Ephemeroptera) in the Oglio River (Azzanello – province of Cremona). *Pianura* 12: 197-200. (In Italian)
- Boonthai, C.; Scott, R. R.; Chapman, R. B. 2000. Acetylcholinesterase as a biomarker to assess the effect of chlorpyrifos and atrazine on some New Zealand aquatic invertebrates. *Australasian J. Ecotoxicol.* 6(1): 59-64.
- Borgman, U.; Norwood, W. P.; Reynoldson, T. B.; Rosa, F. 2001. Identifying cause in sediment assessments: Bioavailability and the Sediment Quality Triad. *Canadian J. Fish. Aquat. Sci.* 58(5): 950-960.
- Bosco, I. J.; Perry, J. A. 2000. Drift and benthic invertebrate responses to stepwise and abrupt increases in non-scouring flow. *Hydrobiologia* 436: 191-208.
- Bouwes, N. W. 1999. The impacts of native and nonnative fish on stream insect communities (Predation). Dissertation, Utah State University.
- Brannstrom, P. A. 1999. Visual ecology of insect superposition eyes. Dissertation, Lunds Universitet.
- Brinkmann, R. 2000. Rediscovery of *Potamanthus luteus* (Ephemeroptera: Potamanthidae) in the Elbe, Saxony-Anhalt. *Entomol. Mitt. Sachsen Anhalt* 8(2): 66. (In German, English summary)
- Burgherr, P. 2001. Spatio-temporal community patterns of lotic zoobenthos across habitat gradients in an alpine glacial stream ecosystem. Dissertation, Eidgenossische Tech. Hochschule Zuerich.
- Callisto, M.; Goulart, M. D. C. 2000. Phoretic association between *Nanocladius* (*Plecopteracoluthus*) sp. (Chironomidae: Diptera) and *Thraulodes* sp. (Leptophlebiidae: Ephemeroptera). *An. Soc. Entomol. Brasil* 29(3): 605-608.
- Callisto, M.; Moreno, P.; Barbosa, F. A. R. 2001. Habitat diversity and benthic functional trophic groups at Serra do Cipo, Southeast Brazil. *Brazilian. J. Biol.* 61(2): 259-266.
- Canivet, V.; Chambon, P.; Gibert, J. 2001. Toxicity and bioaccumulation of arsenic and chromium in epigeal and hypogean freshwater macroinvertebrates. *Arch. Environ. Contam. Toxicol.* 40(3): 345-354.
- Carlisle, D. M. 2001. Trophic structure and function of stream food webs along a gradient of metal contamination. Dissertation, Colorado State Univ.
- Casas, J. J.; Zamora, M. C.; Archila, F.; Alba-Tercedor, J. 2000. The effect of a headwater dam on the use of leaf bags by invertebrate communities. *Regul. Riv. Res. Manage.* 16(6): 577-591.
- Catling, P. M. 2001. Streams and rivers highlighted as major natural area protection priorities in Ontario based on damselfly and dragonfly indicators. *Ontario Odonata* 2: 36-43.
- Chadwick, M. A.; Feminella, J. W. 2001. Influence of salinity and temperature on the growth and production of a freshwater mayfly in the Lower Mobile River, Alabama. *Limnol. Oceanogr.* 46(3): 532-542.
- Chae, J. S.; Pusterla, N.; Johnson, E.; Derock, E.; Lawler, S. P.; Madigan, J. E. 2000. Infection of aquatic insects with trematode metacercariae carrying *Ehrlichia risticii*, the cause of Potomac horse fever. *J. Med. Entomol.* 37(4): 619-625.
- Chambers, P. A.; Brown, S.; Culp, J. M.; Lowell, R. B.; Pietroniro, A. 2000. Dissolved oxygen decline in ice-covered rivers of northern Alberta and its effects on aquatic biota. *J. Aquat. Ecosystem Stress Recovery* 8(1): 27-38.
- Ciborowski, J. J. H.; Corkum L. D.; Krieger, K. A.; Schloesser, D. W.; 1997. Burrowing mayfly (*Hexagenia*) populations in Lake Erie. *Canadian Tech. Rep. Fish. Aquat. Sci.* 2192: 93, [Abstract]
- Cleto, F. S. E. N.; Walker, I. 2001. Effects of urban occupation on the aquatic macroinvertebrates from a small stream of Manaus, Amazonas State, Brazil. *Acta Amazonica* 31(1): 69-89. (In Portuguese; English summary)
- Cooper, B. A. 1999. A physicochemical and benthic macroinvertebrate analysis of three least-impacted first-order streams in east Texas. Dissertation, Stephen F. Austin State University.
- Cox, C. A.; Schultz, R. D.; Guy, C. S. 2001. Diets of white bass in Fall River Reservoir, Kansas. *J. Freshwater Ecol.* 16(3): 429-433.
- Culp, J. M.; Podemski, C. L.; Cash, K. J. 2000. Interactive effects of nutrients and contaminants from pulp mill effluents on riverine benthos. *J. Aquat. Ecosystem*

- Stress Recovery 8(1): 67-75.
- Dean, J. C. 2000. Descriptions of new Leptophlebiidae (Insecta: Ephemeroptera) from Australia. II. *Kaninga*, a new monotypic genus from south-western Australia. *Rec. West. Australian Mus.* 20(1): 87-94.
- Desio, F. 1999. The freshwater macroinvertebrates of Friuli-Venezia Giulia (Northeastern Italy): Mayflies from running waters (Ephemeroptera). *Gortania Atti Mus. Friulano Storia Natur.* 21: 177-201. (In Italian, English abstract)
- de Souza, F. G. M.; Takeda, A. M. 2000. Invertebrates associated with *Paspalum repens* (Poaceae) at the mouth of Caracu Stream (1991-1992), affluent of the Parana River, Porto Rico-PR-Brazil. *Brazilian Arch. Biol. Technol.* 43(3): 317-325.
- Diomande, D.; Gourene, G.; Tito, D. M. L. 2001. Diet of *Synodontis bastiani* (Siluriformes: Mochokidae) Bia River, Ivory Coast. *Cybiurn* 25(1): 7-21.
- Donath, U.; Robinson, C. T. 2001. Ecological characteristics of lake outlets in Alpine environments of the Swiss Alps. *Arch. Hydrobiol.* 150(2): 207-225.
- Doran, B. R. 1999. The macroinvertebrate community of vernal pools in southwestern Quebec. Dissertation, McGill University.
- Drury, D. M.; Kelso, W. E. 2000. Invertebrate colonization of woody debris in coastal plain streams. *Hydrobiologia* 434: 63-72.
- Elouard, J.-M.; Sartori, M. 2001. A revision of the Malagasy genus *Madecocercus* Malzacher, 1995. (Ephemeroptera, Caenidae). *Bull. Soc. Vaudoise Sci. Natur.* 87(3): 229-235.
- Elser, P. 2001. Assessing small-scale directional movements of benthic invertebrates in streams by using a multidirectional cage trap. *Limnologia* 31(2): 119-128.
- Ferreira, R. L.; Horta, L. C. S. 2001. Natural and human impacts on invertebrate communities in Brazilian caves. *Brazilian J. Biol.* 61(1): 7-17.
- Fischer, S.; Marinone, M. C.; Fontanarrosa, M. S.; Nieves, M.; Schweigmann, N. 2000. Urban rain pools: Seasonal dynamics and entomofauna in a park of Buenos Aires. *Hydrobiologia* 441: 45-53.
- Fore, L. S.; Paulsen, K.; O'Laughlin, K. 2001. Assessing the performance of volunteers in monitoring streams. *Freshwater Biol.* 46(1): 109-123.
- Fossati, O.; Wasson, J. G.; Hery, C.; Salinas, G. 2001. Impact of sediment releases on water chemistry and macroinvertebrate communities in clear water Andean streams (Bolivia). *Arch. Hydrobiol.* 151(1): 33-50.
- Francischetti, C. N.; Da Silva, E. R.; Salles, F. F. 2001. On the food of nymphs of *Caenis cuniana* Froehlich, 1969 (Ephemeroptera: Caenidae) in a temporary marsh in Restinga de Marica, Rio de Janeiro State, Brasil. *Bol. Mus. Nac. Rio de Janeiro Zool.* 446: 1-6. (In Portuguese, English summary).
- French, J. R. P., III; Jude, D. J. 2001. Diets and diet overlap of nonindigenous gobies and small benthic native fishes co-inhabiting the St. Clair River, Michigan. *J. Great Lakes Res.* 27(3): 300-311.
- Frost, P. C. 2001. Ecological stoichiometry of trophic interactions in the benthos of boreal lakes. Dissertation, Arizona State University.
- Gaino, E.; Reborá, M. 2000. The duct connecting Malpighian tubules and gut: An ultrastructural and comparative analysis in various Ephemeroptera nymphs (Pterygota). *Zoomorphology* 120(2): 99-106.
- Gaino, E.; Reborá, M. 2001. Synthesis and function of the fibrous layers covering the eggs of *Siphonurus lacustris* (Ephemeroptera, Siphonuridae). *Acta Zool.* 82(2): 41-48.
- Gaino, E.; Sartori, M.; Reborá, M. 2001. Chorionic fine structure of eggs from some species of *Proboscoidoplocia* (Ephemeroptera, Ephemeroidea). *Italian J. Zool.* 68(1): 1-8.
- Galacatos, K. 2001. Community ecology of fishes and aquatic insects in rainforests of the Ecuadorian Amazon. Dissertation, University of New York Coll. Environ. Sci. Forestry.
- Galdean, N. 1997. Some comments about the benthic communities of the Iza River (Maramures, Romania) with special remarks on rheophilic mayfly species (Insecta, Ephemeroptera). *Trav. Mus. Nat. Hist. Natur. Grigore Antipa* 37: 193-203.
- Galdean, N.; Barbosa, F. A. R.; Callisto, M.; Rocha, L. A.; Marques, M. 1999. A proposed typology for the rivers of Serra do Cipo (Minas Gerais, Brazil) based on the diversity of benthic macroinvertebrates and the existing habitats. *Trav. Mus. Nat. Hist. Natur. Grigore Antipa* 41: 445-453.
- Galdean, N.; Callisto, M.; Barbosa, F. A. R. 2001. Biodiversity assessment of benthic macroinvertebrates in altitudinal lotic ecosystems of Serra do Cipo (MG, Brazil). *Brazilian J. Biol.* 61(2): 239-248.
- Gattolliat, J.-L. 2001. Six new species of *Labiobaetis* Novikova & Kluge (Ephemeroptera: Baetidae) from Madagascar with comments on the validity of the genus. *Ann. Limnol.* 37(2): 97-123.
- Gattolliat, J.-L. 2001. *Rheoptilum*: A new genus of two-tailed Baetidae (Ephemeroptera) from Madagascar. *Aquat. Insects* 23(1): 67-81.
- Gattolliat, J.-L. 2001. The genus *Cloeodes* (Ephemeroptera: Baetidae) in Madagascar. *Rev. Suisse Zool.* 108(2): 387-402.
- Gereben, K.; Barbara, A.; Pass, G. 2000. Circulatory organs of abdominal appendages in primitive insects (Hexapoda: Archaeognatha, Zygentoma and Ephemeroptera). *Acta Zool.* 81(4): 285-292.
- Gerlach, J. 2001. The mayflies of Seychelles: Morphology, distribution and ecology. *Phelsuma* 9:67-70.
- Gerlofsma, J. 1999. The effects of anoxia and temperature on the development and survivorship of *Hexagenia* (Ephemeroptera: Ephemeridae) embryos, and implications for western Lake Erie populations. Dissertation, University of Windsor.
- Gewurtz, S. B.; Lazar, R.; Haffner, G. D. 2000. Comparison

- of polycyclic aromatic hydrocarbon and polychlorinated biphenyl dynamics in benthic invertebrates of Lake Erie, USA. *Environ. Toxicol. Chem.* 19(12): 2943-2950.
- Giddings, J. M.; Solomon, K. R.; Maund, S. J. 2001. Probabilistic risk assessment of cotton pyrethroids. II. Aquatic mesocosm and field studies. *Environ. Toxicol. Chem.* 20(3): 660-668.
- Gillies, M. T. 2001. New species of Baetidae (Ephemeroptera) from the Upper River Sigi, Usambara, Tanzania. *J. Natur. Hist.* 35(1): 23-32.
- Godunko, R. J. 2000. Little known species of the genera *Rhithrogena* and *Electrogena* (Ephemeroptera, Heptageniidae) from Ukraine. *Vestn. Zool. Suppl.* 14(1): 60-66.
- Gonzalez, J. M.; Basaguren, A.; Pozo, J. 2001. Life history and production of *Caenis luctuosa* (Burmeister) (Ephemeroptera, Caenidae) in two nearby reaches along a small stream. *Hydrobiologia* 452: 209-215.
- Gorb, S. N.; Kesel, A.; Berger, J. 2000. Microsculpture of the wing surface of Odonata: Evidence for cuticular wax covering. *Arthropod Structure Dev.* 29(2): 129-135.
- Gore, J. A.; Layzer, J. B.; Mead, J. 2001. Macroinvertebrate instream flow studies after 20 years: A role in stream management and restoration. *Regul. Rivers Res. Manage* 17(4-5): 527-542.
- Gualdoni, C. M.; Del C Corigliano, M. 1999. Drift of insects and its structural relationship with the benthos. *Idesia* 17: 57-71. (In Spanish; English summary)
- Haase, P.; Schindehuetten, K. 2000. Ephemeroptera, Plecoptera, aquatic Coleoptera (partim) and Trichoptera from the Lower Saxony part of the Harz Mountains (Germany): Fauna and ecological notes. *Braun. Natur. Schrift.* 6(1): 85-102. (In German; English summary)
- Harrahy, E. A. 2000. Ecotoxicological and genetic effects of a mixture of heavy metals on selected aquatic macroinvertebrates (Colorado). Dissertation, Colorado State University.
- Haybach, A.; Pottgiesser, T. 2000. Mayflies from Northrhine-Westphalia: A first provisional check-list, with new records (Insecta: Ephemeroptera). *Entomol. Z.* 110(11): 322-330. (In German; English summary)
- Hilsenhoff, W. L. 2001. Diversity and classification of insects and Collembola. Pages 661-731. *IN: Thorp, J. H.; Covich, A. P. (eds.). Ecology and classification of North American Freshwater Invertebrates. Second edition. Academic Press, San Francisco.*
- Hilton, D. 2000. It's a small world. Fleeting existence. *Alberta Natur.* 30(2): 38-40.
- Hirasawa, R. 2000. Life history of fish parasitic nematodes in streams. *Aquabiology* 22(5): 424-429. (In Japanese; English summary)
- Hitchings, T. R. 2001. The Canterbury Museum mayfly collection and database (Insecta: Ephemeroptera). *Rec. Canterbury Mus.* 15: 11-32.
- Hoffsten, P. O.; Malmqvist, B. 2000. The macroinvertebrate fauna and hydrogeology of springs in central Sweden. *Hydrobiologia* 436: 91-104.
- Hohmann, M. 2000. First record of *Procloeon pennulatum* (Ephemeroptera: Baetidae) in Saxony-Anhalt. *Entomol. Mitt. Sachsen-Anhalt* 8(2): 67. (In German; English summary)
- Holomuzki, J. R.; Biggs, B. J. F. 2000. Taxon-specific responses to high-flow disturbances in streams: Implications for population persistence. *J. N. Amer. Benthol. Soc.* 19(4): 670-679.
- Hughes, J. M.; Bunn, S. E.; Cleary, C.; Hurwood, D. A. 2000. A hierarchical analysis of the genetic structure of an aquatic insect *Bungona* (Baetidae: Ephemeroptera). *Heredity* 85(6): 561-570.
- Humpesch, U. H.; Elliott, J. M. 1999. Egg hatching: One mechanism for life cycle partitioning in aquatic insects. *Mar. Biol. Ass. United Kingdom Occas. Publ.* 6: 53-66.
- Hurn, A. D.; Riley, R. H.; Young, R. G.; Arbuckle, C. J.; Peacock, K.; Lyon, G. 2001. Temporal shift in contribution of terrestrial organic matter to consumer production in a grassland river. *Freshwater Biol.* 46(2): 213-226.
- Imbert, J. B.; Perry, J. A. 2000. Drift and benthic invertebrate responses to stepwise and abrupt increases in non-scouring flow. *Hydrobiologia* 436: 191-208.
- Inza, B.; Rouleau, C.; Tjalve, H.; Ribeyre, F.; Campbell, P. G. C.; Pelletier, E.; Boudou, A. 2001. Fine scale tissue distribution of cadmium, inorganic mercury, and methylmercury in nymphs of the burrowing mayfly *Hexagenia rigida* studied by whole-body autoradiography. *Environ. Res.* 85(3): 265-271.
- Ishiwata, S. I. 1999. Mayflies of Rishiri-to Island. *Rishiri Stud.* 18: 21-27. (In Japanese)
- Ishiwata, S. I. 2001. Mayflies of Chiba Prefecture, Japan – checklist, diagnoses and keys. *J. Natur. Hist. Mus. Inst. Chiba* 6(2): 163-200. (In Japanese; English summary)
- Ishiwata, S. I.; Tiunova, T. M.; Kuranishi, R. B. 2000. The mayflies (Insecta: Ephemeroptera) collected from the Kamchatka Peninsula and the North Kuril Islands in 1996-1997. *Natur. Hist. Res. Spec. Issue* 7: 67-75.
- Jackson, R. R.; Li, D. 2001. Prey-capture techniques and prey preferences of *Zenodorus durvellei*, *Z. metallescens* and *Z. orbiculatus*, tropical ant-eating jumping spiders (Araneae: Salticidae) from Australia. *New Zealand. J. Zool.* 28(3): 299-341.
- Jacobus, L. M.; McCafferty, W. P. 2001. The mayfly fauna of New York State (Insecta: Ephemeroptera). *J. N. Y. Entomol. Soc.* 109(1): 47-80.
- Jacobus, L. M.; McCafferty, W. P. 2001. Adult descriptions and commentary for two species of southeastern Nearctic *Ephemerella* (Ephemeroptera: Ephemerellidae). *Entomol. News* 112(5): 294-298.
- Jaehnichen, H.; Kahlert, E. 2000. First record of Ephemeroptera, Coptoclava, Coleoptera inc. sed. and Turfanograptus from Lower Cretaceous paper-shales of the western most site of Mongolia. *Mitt. Mus. Natur. Berlin Geowiss.-Reihe* 3: 77-86.
- Jazdzewska, T. 2000. Mayflies (Ephemeroptera) of the Niebieskie Zrodla nature reserve near Tomaszow

- Mazowiecki (central Poland). *Acta Univ. Lodziensis Folia Limnol.* 7: 87-100.
- Johansen, M.; Elliott, J. M.; Klemetsen, A. 2000. Diel fluctuations of invertebrate drift in a Norwegian stream north of the Arctic Circle. *Norwegian J. Entomol.* 47(2): 101-112.
- Kashian, D. R.; Burton, T. M. 2000. A comparison of macroinvertebrates of two Great Lakes coastal wetlands: Testing potential metrics for an index of ecological integrity. *J. Great Lakes Res.* 26(4): 460-481.
- Kelly, Q. M.; Bracken, J. J. 2000. The distribution of Ephemeroptera in Ireland. *Occas. Pub. Irish Biogeogr. Soc.* 5: 1-223.
- Kelly, Q. M.; Bracken, J. J. 2001. Ephemeropteran assemblages in Ireland. *Int. Ver. Theor. Angew. Limnol. Verh.* 27(2): 963-969.
- Kiffney, P. M.; Bull, J. P. 2000. Factors controlling periphyton accrual during summer in headwater streams of southwestern British Columbia, Canada. *J. Freshwater Ecol.* 15(3): 339-351.
- Kiffney, P. M.; Richardson, J. S. 2001. Interactions among nutrients, periphyton, and invertebrate and vertebrate (*Ascapus truei*) grazers in experimental channels. *Copeia* 2001(2): 422-429.
- Kimble, M. S. 1999. Variation of aquatic and terrestrial riparian biodiversity in response to watershed condition. Dissertation, University of Washington.
- Klass, K. D. 2001. The female abdomen of the viviparous earwig *Hemimerus vosseleri* (Insecta: Dermaptera: Hemimeridae), with a discussion of the postgenital abdomen of Insecta. *Zool. J. Linn. Soc.* 131(3): 251-307.
- Klubertanz, T. H.; Hess, C. R. 2001. Additional Illinois and Wisconsin mayfly records, including *Homoeoneuria ammophila* (Ephemeroptera: Oligoneuriidae) in Illinois. *Entomol. News* 112(4): 254, 284.
- Kolligs, D. 2000. Ecological effects of artificial light sources on nocturnally active insects, in particular on butterflies (Lepidoptera). *Faun. Oekol. Mitt. Suppl.* 28: 1-136. (In German; English summary)
- Kovacs, T.; Ambrus, A. 2000. Two rare plecopterans from the Raba: *Agnentina elegantula* (Klapalek, 1905) and *Marthamea vitripennis* (Burmeister, 1839) (Plecoptera: Perlidae). *Misc. Zool. Hungarica* 13: 77-80.
- Kovacs, T.; Ambrus, A.; Juhasz, P. 2000. New Hungarian mayfly (Ephemeroptera) species arising from collectings of larvae II. *Misc. Zool. Hungarica* 13: 81-83.
- Krzeminski, W.; Lombardo, C. 2001. New fossil Ephemeroptera and Coleoptera from the Ladinian (Middle Triassic) of Canton Ticino (Switzerland). *Riv. Itali. Paleontol. Stratigrafia* 107(1): 69-78.
- Kuhara, N.; Nakano, S.; Miyasaka, H. 2000. Flow rate mediates the competitive influence of a grazing caddisfly on mayflies. *Ecol. Res.* 15(2): 145-152.
- Kumar, A.; Dunkel, F. V.; Broughton, M. J.; Sriharan, S. 2000. Effect of root extracts of Mexican marigold, *Tagetes minuta* (Asterales: Asteraceae), on six nontarget aquatic macroinvertebrates. *Environ. Entomol.* 29(2): 140-149.
- Lancaster, J.; Waldron, S. 2001. Stable isotope values of lotic invertebrates: Sources of variation, experimental design, and statistical interpretation. *Limnol. Oceanogr.* 46(3): 723-730.
- Lang, C.; Strawczynski, A.; Vioget, P. 2000. Pesticides and zoobenthic diversity in 23 rivers of western Switzerland: The 1998 and 1999 surveys. *Bull. Soc. Vaudoise Sci. Natur.* 87(2): 93-107. (In French; English summary)
- Leal, J. J. F.; de Assis, E. F. 2000. Life cycle and production of *Campsurus notatus* (Ephemeroptera, Polymitarciidae) in an Amazonian lake impacted by bauxite tailings (Para, Brazil). *Hydrobiologia* 437: 91-99.
- Ledger, M. E.; Winterbourn, M. J. 2000. Growth of New Zealand stream insect larvae in relation to food type. *Arch. Hydrobiol.* 149(3): 353-364.
- Lee, J. O.; Hershey, A. E. 2000. Effects of aquatic bryophytes and long-term fertilization on arctic stream insects. *J. N. Amer. Benthol. Soc.* 19(4): 697-708.
- Lee, J. W.; Quan, Y. T.; Jung, J. C. 2001. Mayflies from Heilongjiang and Jilin Provinces in northeastern China. *Korean J. Entomol.* 31(1): 15-23.
- Lee, S. J.; Bae, Y. J.; Yoon, I. B. 1999. The preference on the substrate particle size and microdistribution of two potamanthid mayflies, *Potamanthus yooni* and *Rhoenanthus coreanus* (Ephemeroptera: Potamanthidae). *Entomol. Res. Bull.* 25: 17-27.
- Lehane, B. M.; Walsh, B.; Giller, P. S.; O'Halloran, J. 2001. The influence of small-scale variation in habitat on winter trout distribution and diet in an afforested catchment. *Aquatic Ecol.* 35(1): 61-71.
- Li, J.; Herlihy, A.; Gerth, W.; Kaufmann, P.; Gregory, S.; Urquhart, S.; Larsen, D. P. 2001. Variability in stream macroinvertebrates at multiple spatial scales. *Freshwater Biol.* 46(1): 87-97.
- Lugo-Ortiz, C. R. 1999. Systematic studies of Baetidae (Insecta: Ephemeroptera), with emphasis on biodiversity in the southern hemisphere. Dissertation, Purdue University.
- Lugo-Ortiz, C. R.; Barber James, H. M.; McCafferty, W. P.; de Moor, F. C. 2001. A non-paraphyletic classification of the Afrotropical genus *Acanthiops* Waltz & McCafferty (Ephemeroptera: Baetidae). *African Entomol.* 9(1): 1-15.
- Lugo-Ortiz, C. R.; de Moor, F. C.; Barber James, H. M. 2000. A taxonomic and ecological review of *Pseudocloeon glaucum* (Agnew) (Ephemeroptera: Baetidae). *African Entomol.* 8(2): 281-288.
- Lugo-Ortiz, C. R.; McCafferty, W. P. 2001. A new species of *Dicentropilum* (Ephemeroptera: Baetidae) from Kenya. *Entomol. News* 112(1): 52-55.
- Lydy, M. J.; Crawford, C. G.; Frey, J. W. 2000. A comparison of selected diversity, similarity, and biotic indices for detecting changes in benthic-invertebrate community structure and stream quality. *Arch. Environ. Contam.*

- Toxicol. 39(4): 469-479.
- Lytle, D. A. 2000. Biotic and abiotic effects of flash flooding in a montane desert stream. *Arch. Hydrobiol.* 150(1): 85-100.
- Lytle, D. A.; Peckarsky, B. L. 2001. Spatial and temporal impacts of a diesel fuel spill on stream invertebrates. *Freshwater Biol.* 46(5): 693-704.
- Maddock, I. P.; Bickerton, M. A.; Spence, R.; Pickering, T. 2001. Reallocation of compensation releases to restore river flows and improve instream habitat availability in the Upper Derwent catchment, Derbyshire, UK. *Regul. Rivers Res. Manage.* 17(4-5): 417-441.
- Mahato, M. 2000. Resource partitioning among larvae of six coexisting odonate species of the Kali Gandaki River, central Nepal (Anisoptera). *Odonatologica* 29(3): 209-223.
- Mahaulpatha, T.; Mahaulpatha, C.; Nakane, K.; Fujii, T. 2001. Nestling diet and prey selection of the Japanese Wagtail *Motacilla grandis*. *J. Yamashina Inst. Ornithol.* 33(1): 36-43.
- Majzlan, O.; Kropilova, B.; Veghova, V. 2000. Colonization of chosen rheolithon groups (Ephemeroptera, Coleoptera, Diptera) on the Bystrica Creek stones in Protected Landscape Area Ponitrie. *Rosalia* 15: 121-130. (In Slovak; English summary)
- Malmqvist, B.; Hoffsten, P. O. 2000. Macroinvertebrate taxonomic richness, community structure and nestedness in Swedish streams. *Arch. Hydrobiol.* 150(1): 29-54.
- Manel, S.; Buckton, S. T.; Ormerod, S. J. 2000. Testing large-scale hypotheses using surveys: The effects of land used on the habitats, invertebrates and birds of Himalayan rivers. *J. Appl. Ecol.* 37(5): 756-770.
- March, J. G. 2000. The role of freshwater shrimps: Patterns and processes along a tropical island stream continuum. Dissertation, University of Georgia.
- Marden, J. H.; O'Donnell, B. C.; Thomas, M. A.; Bye, J. Y. 2000. Surface-skimming stoneflies and mayflies: The taxonomic and mechanical diversity of two-dimensional aerodynamic locomotion. *Physiol. Biochem. Zool.* 73(6): 751-764.
- Margreiter, M.; Kownacki, A.; Kawecka, B.; Kwandrans, J. 1998. Effect of treated wastes on algal and macroinvertebrate communities in an alpine stream. *Oecologia Montana* 7(1-2): 13-20.
- Marklund, O.; Blindow, I. Hargeby, A. 2001. Distribution and diel migration of macroinvertebrates within dense submerged vegetation. *Freshwater Biol.* 46(7): 913-924.
- Marshall, B. D. 2001. An evaluation of the sensitivity of a macroinvertebrate biomonitoring study in headwater streams of New River Gorge National River. *J. Freshwater Ecol.* 16(3): 415-428.
- Marten, M. 2001. Environmental monitoring in Baden-Wuerttemberg with special reference to biocoenotic trend-monitoring of macrozoobenthos in rivers and methodical requirements for evaluation of long-term biocoenotic changes. *Aquatic Ecol.* 35(2): 159-171.
- Masteller, E. C.; Obert, E. C. 2000. Excitement along the shores of Lake Erie – *Hexagenia* – echoes from the past. *Great Lakes Res. Rev.* 5(1): 25-36.
- Mathooko, J. M. 2000. Artificial physical disturbances on streambed sites of a Kenyan mountain stream: Effects of inter-disturbance intervals on mayfly densities. *Limnologica* 30(4): 331-340.
- Mathooko, J. M. 2001. Temporal and spatial distribution of the baetid *Afroptilum sudafricanum* in the sediment surface of a tropical stream. *Hydrobiologia* 443: 1-8.
- Matthaei, C. D.; Townsend, C. R. 2000. Long-term effects of local disturbance history on mobile stream invertebrates. *Oecologia* 125(1): 119-126.
- McCafferty, W. P. 2001. Status of some historically unfamiliar American mayflies (Ephemeroptera). *Pan-Pac. Entomol.* 77(3): 210-218.
- McCafferty, W. P.; Davis, J. R. 2001. Texas Caeninae (Ephemeroptera: Caenidae), with description of a new species. *Proc. Entomol. Soc. Wash.* 103(2): 349-355.
- McCafferty, W. P.; Jacobus, L. M. 2001. Revisions to *Plauditus cestus* and *P. gloveri* (Ephemeroptera: Baetidae). *Entomol. News* 112(5): 305-310.
- McCafferty, W. P.; Klubertanz, T. H.; Randolph, R. P.; Provonsha, A. V.; Lawson, H. R.; Kondratieff, B. C. 2001. Mayflies (Ephemeroptera) of the Great Plains. I: Nebraska. *Trans. Amer. Entomol. Soc.* 127(1): 5-29.
- McEwen, H. A.; Leff, L. G. 2001. Colonization of stream macroinvertebrates by bacteria. *Arch. Hydrobiol.* 151(1): 51-65.
- McKee, D.; Atkinson, D. 2000. The influence of climate change scenarios on populations of the mayfly *Cloeon dipterum*. *Hydrobiologia* 441: 55-62.
- Meadows, D. W. 2001. Effects of bison trampling on stream macroinvertebrate community structure on Antelope Island, Utah. *J. Freshwater Ecol.* 16(1): 83-92.
- Mebane, C. A. 2001. Testing bioassessment metrics: Macroinvertebrate, sculpin, and salmonid responses to stream habitat, sediment, and metals. *Environ. Monit. Assess.* 67(3): 293-322.
- Meyer, M. D.; McCafferty, W. P. 2001. Hagen's small minnow mayfly (Ephemeroptera: Baetidae) in North America. *Entomol. News* 112(4): 255-263.
- Milani, D.; Reynoldson, T. B.; Kirby, S. 1997. Whole sediment toxicity tests and their application in determining the relative sensitivity of four benthic invertebrates to different classes of compounds in spiked and natural sediments. *Canadian Tech. Rep. Fish. Aquat. Sci.* 2192: 33. [Abstract]
- Milner, A. M.; Knudsen, E. E.; Soiseth, C.; Robertson, A. L.; Schell, D.; Phillips, I. T.; Magnusson, K. 2000. Colonization and development of stream communities across a 200-year gradient in Glacier Bay National Park, U.S.A. *Canadian J. Fish. Aquat. Sci.* 57(11): 2319-2335.
- Minshall, G. W.; Royer, T. V.; Robinson, C. T. 2001. Response of the Cache Creek macroinvertebrates during the first 10 years following disturbance by the 1988 Yellowstone wildfires. *Canadian J. Fish. Aquat. Sci.* 58(6): 1077-1088.

- Miserendino, M. L. 2001. Macroinvertebrate assemblages in Andean Patagonian rivers and streams: Environmental relationships. *Hydrobiologia* 444: 147-158.
- Miyasaka, H.; Nakano, S. 2001. Drift dispersal of mayfly nymphs in the presence of chemical and visual cues from diurnal drift and nocturnal benthic foraging fishes. *Freshwater Biol.* 46(9): 1229-1237.
- Molineri, C. 2001. *Traverhyphes*: A new genus of Leptohiphidae for *Leptohiphes indicator* and related species (Insecta, Ephemeroptera). *Spixiana* 24(2): 129-140.
- Monaghan, M. T.; Spaak, P.; Robinson, C. T.; Ward, J. V. 2001. Genetic differentiation of *Baetis alpinus* Pictet (Ephemeroptera: Baetidae). *Heredity* 86(4): 395-403.
- Moravec, F.; Huffman, D. G. 2001. Observations on the biology of *Rhabdochona kidderi texensis*, a parasite of North American cichlids. *J. Helminthol.* 75(2): 197-203.
- Mueller, R.; Schoenfelder, J. 2001. Discovery of *Baetis tracheatus* (Insecta: Ephemeroptera) in Brandenburg. *Lauterbornia* 40: 99-100. (In German; English summary).
- Murphy, T. 2000. Coal tar in Hamilton Harbour: Attempts at biological restoration guidelines. *Environ. Toxicol.* 15(5): 484-495.
- Myers, M. J. 2000. Aquatic invertebrates in island habitats: Dispersal, population genetics, and ecology. Dissertation, University of California, Berkeley.
- Nedeau, E. J. 1999. The effects of an industrial effluent on the quality of a Michigan warmwater stream. Dissertation, Michigan State University.
- Niesler, A. 2000. Estimation of the water quality in Slepjotka stream on the ground of seston and benthos investigations. *Acta Biol. Silesiana* 34(51): 97-118. (In Polish; English summary)
- Nieto, C.; Dominguez, E. 2001. A new species of *Thraulodes* (Ephemeroptera: Leptophlebiidae) from Mexico. *Aquat. Insects* 23(1): 63-66.
- Ogbogu, S. S. 2001. Factors affecting the distribution and abundance of *Cloeon* and *Caenis* (Ephemeroptera) larvae in a tropical impounded river, Nigeria. *African J. Ecol.* 39(1): 106-112.
- Olsen, D. A.; Townsend, C. R.; Matthaei, C. D. 2001. Influence of reach geomorphology on hyporheic communities in a gravel-bed stream. *New Zealand J. Mar. Freshwater Res.* 35(1): 181-190.
- Pagels, J. F.; Smock, L. A.; Sklarew, S. H. 1998. The water shrew, *Sorex palustris* Richardson (Insectivora: Soricidae), and its habitat in Virginia. *Brimleyana* 25: 120-134.
- Peckarsky, B. L.; Taylor, B. W.; Caudill, C. C. 2000. Hydrologic and behavioral constraints on oviposition of stream insects: Implications for adult dispersal. *Oecologia* 125(2): 186-200.
- Peckarsky, B. L.; Taylor, B. W.; McIntosh, A. R.; McPeck, M. A.; Lytle, D. A. 2001. Variation in mayfly size at metamorphosis as a developmental response to risk of predation. *Ecology* 82(3): 740-757.
- Peran, A.; Millan, A.; Velasco, J.; Luisa-Suarez, M. 2001. A method for analyzing indistinguishable overlapping cohorts in insect secondary production studies. *Int. Ver. Theor. Angew. Limnol. Verh.* 27(2): 989-992.
- Peters, W. L.; Peters, J. G. 2000. Discovery of a new genus of Leptophlebiidae: Leptophlebiinae (Ephemeroptera) in Cretaceous amber from New Jersey. Pages 127-131. *IN: Grimaldi, D. (ed.). Studies on fossils in amber, with particular reference to the Cretaceous of New Jersey.* Backhuys Publishers, Leiden.
- Peterson, D. P.; Foote, C. J. 2000. Disturbance of small stream habitat by spawning sockeye salmon in Alaska. *Trans. Amer. Fish. Soc.* 129(4): 924-934.
- Peterson, J. L. 2001. The use of native macroinvertebrates to assess pesticide risk to Oregon streams. Dissertation, Oregon State Univ.
- Petrusso, P. A.; Hayes, D. B. 2001. Invertebrate drift and feeding habits of juvenile chinook salmon in the upper Sacramento River, California. *California Fish Game* 87(1): 1-18.
- Phillips, E. C. 2001. Life history, food habits and production of *Progomphus obscurus* Rambur (Odonata: Gomphidae) in Harmon Creek of east Texas. *Texas J. Sci.* 53(1): 19-28.
- Pires, A. M.; Cowx, I. G.; Coelho, M. M. 2000. Benthic macroinvertebrate communities of intermittent streams in the middle reaches of the Guadiana Basin (Portugal). *Hydrobiologia* 435: 167-175.
- Podemski, C. L. 1999. Ecological effects of a bleached kraft pulp mill effluent on benthic biota of the Athabasca River (*Ameletus subnotatus*, *Baetis tricaudatus*) Alberta. Dissertation, University of Saskatchewan.
- Polegatto, C. M.; Froehlich, C. G. 2001. Functional morphology of the feeding apparatus of the nymph of *Farrodes* sp. (Ephemeroptera: Leptophlebiidae). *Acta Zool.* 82(2): 165-175.
- Pottgiesser, T.; Koch, P.; Sommerhaeuser, M. 1998. Imagos of certain insect groups (Ephemeroptera, Plecoptera, Trichoptera) as indicators for the characterization of waters surrounding lowland brooks. *Verh. Westdeutscher Entomol.* 1997: 187-197. (In German)
- Ramos, E. B. J.; Pino, M. J. M. 2001. Edible insects of Hidalgo, Mexico. *Anal. Inst. Biol. Univ. Nac. Autonoma Mexico Ser. Zool.* 72(1): 43-84. (In Spanish; English summary)
- Randolph, R. P.; McCafferty, W. P. 2001. New species and records of Ephemeroptera (Insecta) from central Mexico. *Dugesiana* 8(1): 15-21.
- Randolph, R. P.; McCafferty, W. P. 2001. Nunavut mayflies (Ephemeroptera): A supplement for far northern North America. *Entomol. News* 112(1): 56-58.
- Reynoldson, T. B.; Rosenberg, D. M.; Resh, V. H. 2001. Comparison of models predicting invertebrate assemblages for biomonitoring in the Fraser River catchment, British Columbia. *Canadian J. Fish. Aquat. Sci.* 58(7): 1395-1410.
- Robert, M.; Cloutier, L. 2001. Summer food habits of

- harlequin ducks in eastern North America. *Wilson Bull.* 113(1): 78-84.
- Royer, T. V.; Robinson, C. T.; Minshall, G. W. 2001. Development of macroinvertebrate-based index for bioassessment of Idaho rivers. *Environ. Manage.* 27(4): 627-636.
- Sadyrin, V. M. 1999. Distribution of Ephemeroptera larvae in water bodies of Republic Komi. *Gidrobiol. Zh.* 35(2): 35-43. (In Russian; English summary)
- Salas, M.; Dudgeon, D. 2001. Stable isotope determination of mayfly (Insecta: Ephemeroptera) food sources in three tropical Asian streams. *Arch. Hydrobiol.* 151(1): 17-32.
- Sander, B.; Allard, P.; Watson, I.; Paine, M.; Guliov, D. 1997. The uses of artificial substrates as a monitoring tool. *Canadian Tech. Rep. Fish. Aquat. Sci.* 2144: 174-181.
- Schloesser, D. W.; Krieger, K. A.; Ciborowski, J. J. H.; Corkum, L. D. 2000. Recolonization and possible recovery of burrowing mayflies (Ephemeroptera: Ephemeridae: *Hexagenia* spp.) in Lake Erie of the Laurentian Great Lakes. *J. Aquat. Ecosystem Stress Recovery* 8(2): 125-141.
- Schloesser, D. W.; Nalepa, T. F. 2001. Changing abundance of *Hexagenia* mayfly nymphs in western Lake Erie of the Laurentian Great Lakes: Impediments to assessment of lake recovery? *Int. Rev. Hydrobiol.* 86(1): 87-103.
- Sharma, B. K.; Sharma, S.; Hussain, Md. 2000. Species composition of zooplankton in a tropical floodplain lake, Assam (N. E. India). *Rec. Zool. Surv. India* 98(3): 85-93.
- Sinitshenkova, N. D. 2000. A review of Triassic mayflies, with a description of new species from Western Siberia and Ukraine (Ephemerida = Ephemeroptera). *Paleontol. J.* 34: S275-S283. [Supplementary issue 3]
- Sinitshenkova, N. D. 2000. New Jersey amber mayflies: The first North American Mesozoic members of the order (Insecta: Ephemeroptera). Pages 111-125. *IN: Grimaldi, D. (ed.). Studies on fossils in amber, with particular reference to the Cretaceous of New Jersey.* Backhuys Publishers, Leiden.
- Smith, G. C. 2000. Rediscovery of the introduced mayfly *Caenis nigropunctata* (Ephemeroptera: Caenidae) in Waimanalo Stream, O'ahu. *Bishop Mus. Occas. Pap.* 64: 30-31.
- Smith, P. J.; Collier, K. J. 2001. Allozyme diversity and population genetic structure of the caddisfly *Orthopsyche fimbriata* and the mayfly *Acanthophlebia cruentata* in New Zealand streams. *Freshwater Biol.* 46(6): 795-805.
- Soldan, T.; Putz, M. 2000. Karyotypes of some Central European mayflies (Ephemeroptera) and their contribution to phylogeny of the order. *Acta Soc. Zool. Bohem.* 64(4): 437-445.
- Soli, A. M. S. 2000. Pesticide and fertilizer contamination of streams adjacent to golf courses and the response of the benthic macroinvertebrate community. Dissertation, University of Maryland College Park.
- Soucek, D. J.; Schmidt, T. S.; Cherry, D. S. 2001. In situ studies with Asian clams (*Corbicula fluminea*) detect acid mine drainage and nutrient inputs in low-order streams. *Canadian J. Fish. Aquat. Sci.* 58(3): 602-608.
- Spacek, J. 1997. Contribution to the fauna of mayflies (Ephemeroptera), stoneflies (Plecoptera) and caddisflies (Trichoptera) in the locality Na Plachte (Eastern Bohemia, Hradec Kralove). *Acta Mus. Regin. Ser. A Sci. Natur.* 25: 89-92. (In Czech; English summary).
- Staicu, G.; Banaduc, D.; Galdean, N. 1998. The structure of some benthic macroinvertebrates and fishes communities in the Viseu watershed, Maramures, Romania. *Trav. Mus. Nat. Hist. Natur. Grigore Antipa* 40: 587-608.
- Suter, P. J.; Pearson, M. J. 2001. Redescription of *Bungona* Harker with new synonyms in the Australian Baetidae (Insecta: Ephemeroptera). *Mem. Mus. Victoria* 58(2): 247-254.
- Tachet, H.; Morse, J. C.; Berly, A. 2001. The larva and pupa of *Pseudoneureclipsis lusitanicus* Malicky, 1980 (Trichoptera: Hydropsychoidea): Description, ecological data and taxonomical considerations. *Aquat. Insects* 23(2): 93-106.
- Takemon, Y. 2000. Reproductive behavior and morphology of *Paraleptophlebia spinosa* (Ephemeroptera: Leptophlebiidae): Implications of variation in copula duration. *Limnology* 1(1): 47-56.
- Tanatmis, M. 2000. The Ephemeroptera (Insecta) fauna of Susurluk (Simav) stream and Manyas lake basin. *Turkiye Entomol. Dergisi* 24(1): 55-67 (In Turkish; English summary)
- Taylor, A. J.; O'Halloran, J. 2001. Diet of Dippers *Cinclus cinclus* during an early winter spate and the possible implications for Dipper populations subjected to climate change. *Bird Study* 48(2): 173-179.
- Taylor, B. W.; McIntosh, A. R.; Peckarsky, B. L. 2001. Sampling stream invertebrates using electroshocking techniques: Implications for basic and applied research. *Canadian J. Fish. Aquat. Sci.* 58(3): 437-445.
- Tonka, T.; Weiser, J. 2000. Iridovirus infection in mayfly larvae. *J. Invertebr. Pathol.* 76(3): 229-231.
- Tseng, M. 2000. Factors affecting life history variation in insects (*Ephemerella subvaria*). Dissertation, University of Toronto.
- Turner, J. W. 2001. A benthic macroinvertebrate analysis of six streams in the Angelina National Forest, Texas. Dissertation, Stephen F. Austin State Univer.
- Tyson, J. T.; Knight, R. L. 2001. Response of yellow perch to changes in the benthic invertebrate community of western Lake Erie. *Trans. Amer. Fish. Soc.* 130(5): 766-782.
- Veliz, M. A. 1999. Relationship of nutrients and mayfly (Ephemeroptera) abundance and diet to periphyton biomass in boreal streams. Dissertation, University of Alberta.
- Victor, T. J.; Reuben, R. 1999. Population dynamics of mosquito immatures and the succession in abundance of

- aquatic insects in rice fields in Madurai, south India. *Indian J. Malariology* 36(1-2): 19-32.
- Wagner, F. H. 2001. Spatial drift distributions and their ecological importance. *Int. Ver. Theor. Angew. Limnol. Verh.* 27(3): 1631-1634.
- Wang, F.; Tessier, A.; Hare, L. 2001. Oxygen measurements in the burrows of freshwater insects. *Freshwater Biol.* 46(3): 317-327.
- Wayland, M.; Hobson, K. A. 2001. Stable carbon, nitrogen, and sulfur isotope ratios in riparian food webs on rivers receiving sewage and pulp-mill effluents. *Canadian J. Zool.* 79(1): 5-15.
- Wellnitz, T. A.; Poff, N. L.; Cosyleon, G.; Steury, B. 2001. Current velocity and spatial scale as determinants of the distribution and abundance of two rheophilic herbivorous insects. *Landscape Ecol.* 16(2): 111-120.
- Wiersema, N. A.; McCafferty, W. P. 2000. Generic revision of the North and Central American Leptohiphidae (Ephemeroptera: Pannota). *Trans. Amer. Entomol. Soc.* 126(3-4): 337-371.
- Wiersema, N. A.; W. P. McCafferty; Baumgardner, D. E. 2001. *Asioplax numinuh*, a new species of Ephemeroptera (Leptohiphidae) from Texas and Mexico. *Entomol. News* 112(5): 301-304.
- Williams, J. K.; Townsend, C. R.; Poulin, R. 2001. Mermithid nematode infections and drift in the mayfly *Deleatidium* spp. (Ephemeroptera). *J. Parasitol.* 87(5): 1225-1227.
- Wright, A. B.; Smock, L. A. 2001. Macroinvertebrate community structure and production in a low-gradient stream in an undisturbed watershed. *Arch. Hydrobiol.* 152(2): 297-313.
- Wright, J. F.; Gunn, R. J. M.; Blackburn, J. H.; Grieve, N. J.; Winder, J. M.; Davy, B. J. 2000. Macroinvertebrate frequency data for the RIVPACS III sites in Northern Ireland and some comparisons with equivalent data for Great Britain. *Aquat. Conserv.* 10(5): 371-389.
- Zahradkova, S. 1999. Ephemeroptera. *Folia Fac. Sci. Natur. Univ. Masarykianae Brun. Biol.* 101: 167-171.
- Zajdlik, B. A.; Moran, T.; Munro, S. 1997. Assessing spatial extent of impacted areas in the St. Clair River using the sediment quality triad. *Canadian Tech. Rep. Fish. Aquat. Sci.* 2192: 30. [Abstract]
- Zhou, C. F.; Gui, H.; Su, C. R. 2000. The first record of the genus *Brachycercus* in China with description of a new species (Ephemeroptera: Caenidae). *Entomol. Sinica* 7(2): 132-134.
- Zhou, C. F.; Su, C. R.; Gui, H. 2000. The first record of the genus *Torleya* in China with description of a new species (Ephemeroptera: Ephemerellidae). *Acta Zootaxonomica Sinica* 25(3): 312-314. (In Chinese: English summary)
- Zieba, J. 1996. Chironomidae in the benthos of the submontane Roznow and Czchow Reservoirs (the River Dunajec, southern Poland) 40 years after filling. *Acta Hydrobiol.* 38(3-4): 133-144.

SWOSU

The Mayfly Newsletter (ISSN 1091-4935) is the official newsletter of the International Conferences on Ephemeroptera and is published twice a year to facilitate communication among ephemeropterists. Subscriptions to the *Newsletter* are free. To place your name on the mailing list or to contribute information for the next issue, contact Peter M. Grant, editor, *The Mayfly Newsletter*, Department of Biological Sciences, Southwestern Oklahoma State University, 100 Campus Drive, Weatherford, Oklahoma 73096-3098 USA, phone (580) 774-3294, FAX (580) 774-7140, email grantp@swosu.edu. This publication was authorized by the Dean of Arts and Sciences and was printed at a cost of \$375.00 for 500 copies.

**Permission was not obtained
to publish cartoon online.**



Address Updates

Corrections or additions to the information in the mailing list since the last issue are listed below. Updated addresses will be published as they become available. Please inform the editor of any changes in postal or email addresses.

K ADKINS
1517 WINDWARD LANE
WYLIE TX 75098
USA
allseason@dallas.net

N ARUN NAGENDRAN
DEPARTMENT OF ZOOLOGY AND
MICROBIOLOGY
THIAGARAJAR COLLEGE
MADURAI 625 009 TAMIL NADU
INDIA
naga_arun@yahoo.co.in

B BOONSOONG
DEPARTMENT OF BIOLOGY
FACULTY OF SCIENCE
KHON KAEN UNIVERSITY
KHON KAEN 40002
THAILAND
bboonsoong@yahoo.com

E BRIGHT
2110 INDEPENDENCE BOULEVARD
ANN ARBOR MI 48104-6439
USA
ethanbr@umich.edu

J H CARABELLO
CPRi COMMUNICATIONS INC
370 NORTH STREET
TETERBORO NJ 07608
USA
JCARABELLO@cpronline.com

T FOSTER
4652 SHAY TERRACE
BUFORD GA 30519
USA
TrishAFoster@aol.com

L GRAY
DEPT BIOLOGY
UTAH VALLEY STATE COLLEGE
800 WEST UNIVERSITY PARKWAY
OREM UT 84058-0001
USA
GRAYLA@uvsc.edu

U H HUMPEsch
LIMNOLOGICAL INSTITUTE
MONDSEESTRAßE 9
A 5310 MONDSEE
AUSTRIA
Uwe.Humpesch@oeaw.ac.at

W S LESKO
NORTH JERSEY EYE ASSOCIATES PA
1005 CLIFTON AVENUE
CLIFTON NJ 07013
USA
dryflymd@aol.com

G MARKIEWICZ
DETROIT EDISON-FERMI 2
6400 NORTH DIXIE HIGHWAY 180 TAC
NEWPORT MI 49230
USA
markiewiczg@dteenergy.com

T H OGDEN
DEPARTMENT OF ZOOLOGY
574 WIDB
BRIGHAM YOUNG UNIVERSITY
PROVO UT 84602-5255
USA
heath_ogden@byu.edu

H REUSCH
WELLENDORF 30
D-29562 SUHLENDORF
GERMANY
Herbert.Reusch@t-online.de

A B RICHARDS
CA DEPT FISH AND GAME
AQUAT BIOASSESSMENT LAB
COLLEGE OF NATURAL SCIENCES
CSU-CHICO
CHICO CA 95929-0555
USA
batrichards@earthlink.net

F F SALLES
MUSEU DE ENTOMOLOGIA DEPT
BIOLOGIA ANIMAL
UNIVERSIDADE FEDERAL DE VICOSA
VICOSA MINAS GERAIS CEP 36 571-000
BRASIL
ffsalles@insecta.ufv.br

E SANAA
HOVSGOL GEF PROJECT GEOECOLOGY
INSTITUTE
POB 81 BARUUN SELBE 15
ULAANBAATAR 211238
MONGOLIA
sanaaspu@yahoo.com

H M SAVAGE
HMS1@CDC.GOV

J SLUSARK
CA DEPT FISH AND GAME
AQUAT BIOASSESSMENT LAB
COLLEGE OF NATURAL SCIENCES
CSU-CHICO
CHICO CA 95929-0555
USA

D STALLWOOD
PO BOX 211
WOOD RIVER NE 68883-0211
USA

A H STANICZEK
STAAT MUS NATURKUNDE STUTTGART
ABT PALAEONTOL SEKT BERNSTEIN
ROSENSTEIN 1
D-70191 STUTTGART
GERMANY
staniczek.smns@naturkundemuseum-bw.de

A STAUDER
CARL-JUSTI-STR- 22
53121 BONN
GERMANY
AngelikaStauder@web.de

J M STEINER
BLUESTEM INC
28340 KYSTER ROAD
MADISON WI 54856-2074
USA

D G STROM
PO BOX 7130
PORT ST LUCIE FL 34985-7130
USA
7723321915@mobile.mycingular.net

T TSUTSUMI
FACULTY OF EDUCATION
FUKUSHIMA UNIVERSITY
1 KANAYAGAWA
FUKUSHIMA 960-1296
JAPAN
tsutsumi@educ.fukushima-u.ac.jp

C ZHOU
cfzhou@eyou.com

Unknown Address

P HUBBELL
1471 SOUTHWOOD DRIVE
HUNTSVILLE TX 77340 USA